

Champion Developer

Complete Diet For Beef Cattle.

MEDICATED

For the prevention and control of coccidiosis due to *Eimeria bovis* and *Eimeria zuernii* in cattle being feed in confinement for slaughter.

ACTIVE DRUG INGREDIENT

Monensin 37.0 g/ton

GUARANTEED ANALYSIS

Crude Protein	(Min)	12.00 %
Crude Fat	(MIN)	4.00 %
Crude Fiber	(Max)	4.00 %
Calcium	(Min) 0.60 (Max)	0.75 %
Phosphorus	(Min)	0.30 %
Salt	(Min) 0.35 (Max)	0.50 %
Potassium	(Min)	0.50 %
Magnesium		0 %
Zinc		ppm
Copper	(Min)	35.0 ppm
Selenium	(Min)	0.25 ppm



Vitamin A  **Diamond V** ¹lb
Vitamin E (1000) 0.010/lb

LIST OF INGREDIENTS

Grain Products, Plant Protein Products, Roughage Products, Molasses, Animal Fat, Calcium Carbonate, Salt, Potassium Chloride, Yeast Culture, Magnesium Oxide, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Chromium Propionate, Artificial Flavor, Copper Sulfate, Sodium Selenite, Vitamin E Supplement, Vitamin E Supplement, Vitamin D3 Supplement, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydriodide, Calcium Iodinate, Cobalt Sulfate, Cobalt Carbonate, Mold Max.

FEEDING DIRECTIONS

Consult your Spencer Ag feed consultant for feeding directions for Champion Developer. Provide salt and mineral free choice. Provide an abundant clean, fresh source of water.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use.

Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. CAUTION: This feed contains added copper, do not feed to sheep. CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

Manufactured for



Spencer, IA 51301
(800) 543-8854

No warranty of results is made due to differences in environment, genetic, health, and management.

10/6/2017

NET WT 50 LB (22.68 kg) or BULK

Code No.: D17277